



**Faculty of Resource Science and Technology**

**Detection and Molecular Characterization of *Bacillus cereus* Isolated  
from Sago Processing Plants in Sarawak**

**Jasmin binti Jaraee**

**Master of Science  
2018**

Grade: \_\_\_\_\_

Please tick (✓)

Final Year Project Report

Masters

PhD

<input type="checkbox"/>
<input checked="" type="checkbox"/>
<input type="checkbox"/>

### DECLARATION OF ORIGINAL WORK

This declaration is made on the .....15.....day of ~~SEPTEMBER~~ August 2018.

#### Student's Declaration:

I JASMIN BINTI JARA'EE, 1502198, FACULTY OF RESOURCE SCIENCE AND TECHNOLOGY

(PLEASE INDICATE STUDENT'S NAME, MATRIC NO. AND FACULTY) hereby declare that the work entitled, DETECTION AND MOLECULAR CHARACTERISATION OF BACILLUS CEREUS IN SAGO PROCESSING PLANTS IN SARAWAK is my original work. I have not copied from any other students' work or from any other sources except where due reference or acknowledgement is made explicitly in the text, nor has any part been written for me by another person.

28/9/2018

Date submitted

JASMIN BINTI JARA'EE (15020198)

Name of the student (Matric No.)

#### Supervisor's Declaration:

I LESLIE MAURICE BILUNG (SUPERVISOR'S NAME) hereby certifies that the work entitled, DETECTION AND MOLECULAR CHARACTERISATION OF BACILLUS CEREUS IN SAGO PROCESSING PLANTS IN SARAWAK (TITLE) was prepared by the above named student, and was submitted to the "FACULTY" as a \* partial/full fulfillment for the conferment of MASTER OF SCIENCE (MICROBIOLOGY) (PLEASE INDICATE THE DEGREE), and the aforementioned work, to the best of my knowledge, is the said student's work

Received for examination by: LESLIE MAURICE BILUNG  
(Name of the supervisor)

Date: 26/9/18

I declare this Project/Thesis is classified as (Please tick (✓)):

- ☐ **CONFIDENTIAL** (Contains confidential information under the Official Secret Act 1972)\*  
☐ **RESTRICTED** (Contains restricted information as specified by the organisation where research was done)\*  
☒ **OPEN ACCESS**

#### Validation of Project/Thesis

I therefore duly affirmed with free consent and willingness declared that this said Project/Thesis shall be placed officially in the Centre for Academic Information Services with the abide interest and rights as follows:

- This Project/Thesis is the sole legal property of Universiti Malaysia Sarawak (UNIMAS).
- The Centre for Academic Information Services has the lawful right to make copies for the purpose of academic and research only and not for other purpose.
- The Centre for Academic Information Services has the lawful right to digitise the content to for the Local Content Database.
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis for academic exchange between Higher Learning Institute.
- No dispute or any claim shall arise from the student itself neither third party on this Project/Thesis once it becomes sole property of UNIMAS.
- This Project/Thesis or any material, data and information related to it shall not be distributed, published or disclosed to any party by the student except with UNIMAS permission.

Student's signature \_\_\_\_\_



27.9.2018

(Date)

Supervisor's signature: \_\_\_\_\_



26.9.2018

Dr (Date) Maurice Hilung  
Penyelaras Lantikan Industri  
Program Biologi/Prologi Sains  
Fakulti Sains dan Teknologi Sains  
Universiti Malaysia Sarawak

Current Address:

132 A, KAMPUNG PANGLIMA SEMAN ULU, 93050, KUCHING SARAWAK.

Notes: \* If the Project/Thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach together as annexure a letter from the organisation with the period and reasons of confidentiality and restriction.

[The instrument was duly prepared by The Centre for Academic Information Services]

Detection and Molecular Characterization of *Bacillus cereus* Isolated from Sago  
Processing Plants in Sarawak

Jasmin binti Jaraee

A thesis submitted

In fulfilment of the requirements for the degree of Master of Science (Microbiology)

Faculty of Resource Science and Technology  
UNIVERSITI MALAYSIA SARAWAK  
2018

## **DECLARATION**

The thesis has not been accepted for any degree and is not concurrently submitted in candidature for any other degree.

---

Name: Jasmin binti Jaraee

Matric number: 15020198

30 January 2018

## **ACKNOWLEDGEMENT**

All praises to Allah the Almighty for the blessings, strength and love He has given to me and for granting me the strength, determination and patience to accomplish my master degree. A special thanks and sincere appreciation I dedicated to my respected supervisor, Dr. Lesley Maurice Bilung and my co-supervisors, Assoc. Prof. Dr. Cirilo Hipolito Nolasco and Dr. Micky Vincent for their advices, guidance, encouragement as well as skills and knowledge sharing throughout my study. Deepest gratitude and a big thank you to my family who had supported me morally as well as financially in time of difficulties. I am also deeply indebted to the members of Microbiology Laboratory especially Pui Chai Fung, Jennifer Jalan, Ahmad Syatir Mohd Tahar, Auns Mohamed and Grace Bebey for all their help, moral support and encouragement along my study journey. Not to forget, the lab assistants and UNIMAS driver for their help and service especially during sampling trips. Last but not least, thank you to everyone who helped and supported me directly and indirectly. Without your support, I would never be able to finish this project.

## ABSTRACT

Sago processing industries are well-established and high potential industries in Sarawak. However, the contamination of bacteria might deteriorate its quality and become a concern to the public health. This study aimed to detect, quantify and characterize *Bacillus cereus* in sago processing in Sarawak, Malaysia. *B. cereus* was isolated from two selected sago processing mills in Sarawak. The prevalence and concentration of *B. cereus* in this study were determined firstly using selective agar and followed by using specific Polymerase Chain Reaction (PCR) by targeting specific virulence gene, haemolysin (*hly*) gene. A total of 120 samples consist of bark swab, sago pith, starch slurry, sago milk, sago flour and sago effluent were collected from each processing step in sago mills. It was revealed that *B. cereus* were present in 35% (42/120) of the samples. These isolates were subjected to molecular typing by using Enterobacterial Repetitive Intergenic Consensus Polymerase Chain Reaction (ERIC PCR) and Pulsed Field Gel Electrophoresis (PFGE). Both molecular typing method showed heterogeneity of the *B. cereus* strain. Susceptibility of all isolates towards 14 antibiotics was assessed using disk diffusion assay. *B. cereus* isolates were uniformly resistant to penicillin and ampicillin whereas *B. cereus* isolates were uniformly susceptible to imipenem and norfloxacin. Multiple antibiotic resistance (MAR) index were calculated based on the antibiotic resistance results. The MAR index for all isolates were also varies, ranged from 0.083 to 0.750. This study is useful in developing appropriate intervention strategies and establishing food safety standards in sago processing in Sarawak thus contribute in lowering the disease burden and assist in providing safer food to the society.

**Keywords:** *B. cereus*, sago processing, detection, polymerase chain reaction, molecular characterization, antibiotic susceptibilities test

***Pengesanan dan Pencirian Molekul Bacillus cereus yang Diasingkan dalam Kilang  
Pemprosesan Sagu di Sarawak***

**ABSTRAK**

*Industri pemprosesan sagu merupakan industri yang dikenali dan berpotensi tinggi di Sarawak. Walau bagaimanapun, pencemaran bakteria akan mengurangkan kualitinya dan menjadi isu kepada kesihatan awam. Kajian ini bertujuan untuk mengesan, mengira dan mencirikan B. cereus dalam pemprosesan sagu di Sarawak. B. cereus diasingkan daripada dua kilang pemprosesan sagu yang terpilih di Sarawak. Kelaziman dan penumpuan B. cereus dalam kajian ini dikaji dengan menggunakan agar selektif dan kemudian diuji dengan reaksi polimeras berantai spesifik untuk mengekspresikan gen sasaran iaitu gen hemolaisin. Sejumlah 120 sampel yang terdiri daripada olesan batang, empulur sagu, buburan kanji, susu sagu, tepung sagu dan efluen sagu diperolehi daripada setiap peringkat pemprosesan kedua-dua kilang. Ia menunjukkan bahawa B. cereus wujud di dalam 35% (42/120) sampel. Isolat ini tertakluk kepada pencirian molekul dengan menggunakan ERIC-PCR dan PFGE. Hasil daripada kedua-dua pencirian molekul ini menunjukkan kepelbagaian jenis B. cereus. Sensitiviti kesemua isolat terhadap 14 antibiotik ditaksir menggunakan B. cereus menunjukkan kerintangan kepada penisilin dan ampicilin serta sensitif kepada imipenem dan norfloxazin. Kajian ini berguna dalam membangunkan strategi untuk langkah pembaikan serta menghasilkan piawaian keselamatan makanan dalam pemprosesan sagu di Sarawak sekaligus menyumbang dalam mengurangkan beban penyakit dan membantu ke arah penyediaan makanan yang selamat kepada masyarakat.*

***Kata kunci:*** B. cereus, pemprosesan sagu, pengesanan, tindak balas polimeras berantai, pencirian molekul, ujian sensitiviti antibiotik.



## TABLE OF CONTENTS

	Page
<b>DECLARATION</b>	i
<b>ACKNOWLEDGEMENT</b>	ii
<b>ABSTRACT</b>	iii
<b><i>ABSTRAK</i></b>	iv
<b>TABLE OF CONTENTS</b>	v
<b>LIST OF TABLES</b>	viii
<b>LIST OF FIGURES</b>	ix
<b>LIST OF ABBREVIATIONS</b>	xi
<b>CHAPTER 1: INTRODUCTION AND LITERATURE REVIEW</b>	1
1.1 Introduction and objectives	1
1.2 Literature review	4
1.2.1 Sago palm ( <i>Metroxylon sagu</i> )	4
1.2.2 Sago processing	5
1.2.3 Morphology and characteristics of <i>B. cereus</i>	8
1.2.4 Toxin gene of <i>B. cereus</i>	10
1.2.5 Outbreaks of <i>B. cereus</i>	11
1.2.6 Prevalence of <i>B. cereus</i> in foods and food processing plants	12
1.2.7 Isolation and enumeration of <i>B. cereus</i> by using conventional method	13
1.2.8 Isolation and enumeration of <i>B. cereus</i> by using polymerase chain reaction (PCR)	15

1.2.9	Molecular typing of <i>B. cereus</i>	17
1.2.10	Antibiotic susceptibilities test of <i>B. cereus</i>	19
1.2.11	Multi-antibiotic resistance index	20
<b>CHAPTER 2: MATERIALS AND METHODS</b>		<b>23</b>
2.1	Sample collection	23
2.2	Sample processing	24
2.3	Isolation of <i>B. cereus</i>	24
2.4	Genomic DNA extraction	25
2.5	Detection of haemolysin ( <i>hly</i> ) gene by using PCR	26
2.6	Antibiotic susceptibility test (AST)	28
2.7	Molecular typing of <i>B. cereus</i> associated with sago processing in Sarawak	29
2.7.1	ERIC-PCR	29
2.7.2	PFGE	29
<b>CHAPTER 3: RESULTS</b>		<b>32</b>
3.1	Prevalence of <i>B. cereus</i> in sago-related samples associated with sago processing in Sarawak	32
3.1.1	Identification of <i>B. cereus</i> by polymerase chain reaction (PCR)	32
3.2	Antibiotic Susceptibility Test	35
3.3	Molecular typing of <i>B. cereus</i> associated with sago processing in Sarawak.	46
3.2.1	ERIC PCR	46
3.2.2	PFGE	53

<b>CHAPTER 4: DISCUSSION</b>	59
4.1 Prevalence of <i>B. cereus</i> associated with sago processing in Sarawak	59
4.2 Antibiotic susceptibitiliy test	63
4.3 Molecular typing of <i>B. cereus</i> associated with sago processing in Sarawak	66
<b>CHAPTER 5: CONCLUSION</b>	70
<b>REFERENCES</b>	72
<b>APPENDICES</b>	88

## LIST OF TABLE

	<b>Page</b>
Table 2.1     The primers used	26
Table 2.2     The PCR components	27
Table 2.3     Cycling conditions for haemolysin gene	27
Table 2.4     Condition of ERIC-PCR of <i>B. cereus</i>	29
Table 2.5     Restriction mixture of each microtube	31
Table 2.6     Gel electrophoresis running parameter	31
Table 3.1     The prevalence of <i>B. cereus</i> from sago processing A and sago processing B based on PCR assay	33
Table 3.2     Percentage of bacterial resistance of all <i>B. cereus</i> isolated based on antibiotics	35
Table 3.3     Percentage of antibiotic resistance of <i>B. cereus</i> isolated at sago processing mill A	37
Table 3.4     Percentage of antibiotic resistance of <i>B. cereus</i> isolated at sago processing mill B	38
Table 3.5     Antibigram of <i>B. cereus</i> in sago processing in Sarawak	44
Table 3.6     Antibiotic resistance, ERIC-PCR and PFGE pattern of <i>B. cereus</i> from sago processing in Sarawak	57

## LIST OF FIGURES

	<b>Page</b>
Figure 1.1	5
Figure 1.2	6
Figure 1.3	7
Figure 1.4	7
Figure 1.5	8
Figure 1.6	8
Figure 1.7	9
Figure 1.8	16
Figure 2.1	23
Figure 3.1	32

Figure 3.2	Percentage of antibiotic resistancy-susceptibility of <i>B. cereus</i> in sago processing in Sarawak.	36
Figure 3.3	The percentage of antibiotic resistance of <i>B. cereus</i> in every stage of processing in sago processing mill A.	40
Figure 3.4	The percentage of antibiotic resistance of <i>B. cereus</i> in every stage of processing in sago processing mill B.	41
Figure 3.5	MAR index of <i>B. cereus</i> in sago processing in Sarawak.	46
Figure 3.6	Banding profiles of ERIC-PCR for sago processing A and B.	48
Figure 3.7	Banding profiles of ERIC-PCR for sago processing B.	49
Figure 3.8	Banding profiles of ERIC-PCR for sago processing B.	50
Figure 3.9	Dendrogram of ERIC-PCR profiles for sago processing mill A	51
Figure 3.10	Dendrogram of ERIC-PCR profiles for sago processing mill B	52
Figure 3.11	Banding profiles of PFGE for sago processing A.	54
Figure 3.12	Banding profiles of PFGE for sago processing B.	55
Figure 3.13	Dendrogram of PFGE profiles for sago processing mill A and B.	56

## LIST OF ABBREVIATIONS

μL	Microliter
μm	Micrometer
°C	Degree Celsius
%	Percent
AFLP	Amplified fragment length polymorphism
AGE	Agarose gel electrophoresis
AOAC	Association of Official Analytical Chemists
AST	Antibiotic susceptibility test
CDC	Centre for Disease Control
CFU	Colony forming unit
ddH <sub>2</sub> O	Double distilled water
dNTP	Deoxynucleotide triphosphate
DNA	Deoxyribonucleic acid
EDTA	Ethylenediamine tetra-acetic acid
ERIC	Enterobacteria repetitive intergenic consensus sequences
EtBr	Ethidium bromide
FDA	Food and Drug Association
FSIS	Food Safety and Inspection Service
ISO	International Organization of Standardization
HBL	Haemolysin BL
kb	Kilo base
kDa	Kilo Dalton

MAR	Multiple antibiotic resistant
MgCl <sub>2</sub>	Magnesium Chloride
mL	Milliliter
MLST	Multilocus sequence typing
Mm	Milimeter
mM	Milimolar
MPN	Most probable number
MSRV	Modified Semisolid Rappaport-Vassiliades
MYP	Mannitol-Egg Yolk-Polymyxin
NCBI	National Center for Biotechnology Information
n.d	No date
NGFIS	Netherlands Government Food Inspection Services
NHE	non-hemolytic enterotoxin
NMKL	Nordic Committee on Food Analysis
PCR	Polymerase chain reaction
PEMBA	Polymyxin pyruvate egg-yolk mannitol–bromothymol blue agar
PFGE	Pulsed-field gel electrophoresis
RAPD	Random amplification of polymorphic DNA
rep-PCR	Repetitive Polymerase Chain Reaction
RM	Ringgit Malaysia
Rpm	Revolution per minute
RTE	Ready-to-eat
SLST	Single locus sequence typing



U	Unit
UV	Ultra violet
<i>Taq</i>	<i>Thermus aquaticus</i>
TBE	Tris-borate-EDTA
TBS	Tryptone soy broth
USDA	United States Department of Agriculture
VNTR	Variable number tandem repeat
WHO	World Health Organization



## **CHAPTER 1**

### **INTRODUCTION AND LITERATURE REVIEW**

#### **1.1 Introduction**

In Sarawak, sago processing industries is a well-established industry that contribute to Sarawak export revenue (Karim *et al.*, 2008). Sago palm is the plant that has high potential agricultural industry and is grown commercially in Southern East Asia countries (Singhal *et al.*, 2008). According to Adeni *et al.* (2009), sago is very important food product for over a million peoples as their primary dietary starch source. Starch is one of the most abundant plant products in the world. It is a major source of energy in human daily food. Every year, the consumption of the sago starch by the global citizen is estimated to be between 200,000 to 300,000 tons (Bujang and Muniandy, 2004). The export of sago product from Sarawak in 2013 was estimated to be 50,000 tonne which procuring income approximately RM 81 millions (Department of Agriculture Sarawak, 2013). Usually, sago starch is used for various applications in Malaysia such as glucose, monosodium glutamate and noodles (Bujang, 2008).

##### **1.1.1 Problem Statement**

In food processing industries, some of the preparation, processing and storage procedures were exposed to the risk of bacterial contamination. It might harbor a great number of microorganisms that may cause the food products to spoil and represents a direct health hazards to consumers (Lesley *et al.*, 2013). Most of the incidence of diarrhoea in developing countries are caused by foodborne and waterborne pathogens (WHO, 2008). Diarrhoeal diseases are included in 20 leading causes of death in the world in 2014 (WHO, 2014). Unhygienic food handling and preparation and inadequate cooking which may lead to food

poisoning due to the presence of bacteria. (Sandra *et al.*, 2012). Until March 2015, 143 cases of food poisoning were reported in Sarawak (Sarawak EPID Health News, 2015). There are a few food poisoning bacteria reported to be commonly found in food and its processing which include *Salmonella* spp., *B. cereus*, *Escherichia coli* and *Listeria* species (Greenhill *et al.*, 2007).

*B. cereus* is chosen as potential hazard due to its ubiquitous nature and its preference to live in soil and starchy food. The presence of the *B. cereus* in food processing industries cause problems by reducing the quality of the products (Lesley *et al.*, 2013; Sandra *et al.*, 2012), and affecting people's health after eating the contaminated foods (Ghelardi *et al.*, 2002). *B. cereus* was first discovered by Frankland *et al.* (1887) in the air of the cowshed. *B. cereus* affects foods processing industry both by reducing the quality of the products (Lesley *et al.*, 2013; Sandra *et al.*, 2012) and by endangering people's health upon eating contaminated foods (Ghelardi *et al.*, 2002).

However, to date, there is limited study of the occurrence of microorganisms including *B. cereus* in sago-associated products has been conducted (Greenhill *et al.*, 2007). To the best of our knowledge, there has been no study on the prevalence of *B. cereus* in sago processing has been reported in Sarawak. Hence, in our study we isolated and identified *B. cereus* from sago processing and determine the prevalence of the *B. cereus*. There are several international standard protocol commonly used for isolation and identification of *B. cereus* which include the US Food and Drug Administration (FDA), the Association of Official Analytical Chemists (AOAC), the European Committee for Standardisation (CEN, EN), the International Organization for Standardization (ISO), the Netherlands Government Food Inspection Services (NGFIS), the Nordic Committee on Food Analysis (NMKL) and the United States Department of Agriculture (USDA) Food Safety and Inspection Service (FSIS). In our study, we applied

FDA method which involve the sample enrichment by using tryptone soy broth and direct plating on MYP agar (Lesley *et al.*, 2013).

*B. cereus* is further identified and confirmed by PCR assay. Further confirmation and identification by PCR assay were required as conventional detecting method often give false-positive/false-negative results. Hence PCR assay can be applied as it is reliable, specific, sensitive and reproducible method. In the present study, we detect *B. cereus* by targeting haemolysin gene as described by Mutashar *et al.* (2015), Fukushima *et al.* (2003) and Wang *et al.* (1997). The raw material quality and the processing environment of food products can affect the final products quality (Oh *et al.*, 2012). As contamination and/or cross-contamination were the major concern in food processing, hence we decided to conduct molecular typing of *B. cereus* by using rep-PCR method in order to determine the contamination pattern in the sago processing mills. It is important to understand the routes of contamination of *B. cereus* in sago processing to prevent the contamination of the final products (Oh *et al.*, 2012).

*B. cereus* is an opportunistic pathogen that can cause severe foodborne illness. There were also several studies on the resistance of *B. cereus* toward antibiotics had been reported (Yim *et al.*, 2015; Lee *et al.*, 2012; Park *et al.*, 2009; Rosenquist *et al.*, 2005 and Andrews *et al.*, 2002). This showed it is necessary to investigate the *B. cereus* antibiotic resistance and susceptibility. Antibiotics have been widely used in agriculture, veterinary and medicine. Prolonged usage and exposure towards commonly used antibiotics including penicillin, ampicillin, norfloxazin, imipenem, tetracycline, chloramphenicol, quinolones, aminoglycoside, spectinomycin, cephalosporin, nitrofurantoin, nitroimidazole, sulfonamide, and trimethoprim are able to promote the emergence of resistance in bacteria. It is important to monitor antibiotic susceptibility of *B. cereus* in food processing to ensure and monitor the emergence and spread of bacterial resistance to antimicrobial agents. However, there is no data of antibiotic resistance

of bacteria associated with sago processing mills. In this study, the antibiotic susceptibility of *B. cereus* in the sago was examined by disk diffusion method (Park *et al.*, 2009). The findings of this study provided baseline data on *B. cereus* contamination for future risk assessment work and establishment of food safety standards in sago processing in Sarawak.

### 1.1.2 Objectives

The main goal of this study was to detect, quantify and characterize *B. cereus* in sago processing in Sarawak, Malaysia. The specific objectives of this study were to:

- a. Detect *B. cereus* in sago processing by using Polymerase Chain Reaction.
- b. Determine the contamination level of *B. cereus* in sago processing by enumerating *B. cereus* using standard plate count method.
- c. Determine the antibiotic resistance profiles of *B. cereus* isolated from sago processing.
- d. Characterise *B. cereus* in sago processing by using Repetitive Intergenic Consensus Polymerase Chain Reaction (ERIC- PCR) and Pulsed-field Gel Electrophoresis (PFGE).

## 1.2 Literature review

### 1.2.1 Sago palm

Sago palm (Figure 1.1) or *Metroxylon sagu* belongs to the genus *Metroxylon* of Palmae family (Greenhill *et al.*, 2007). *Metroxylon sagu* is derived from the term ‘metra’ which means pith or parenchyma and the term ‘xylon’ means xylem. Many scientists considered it as the starch crop of the 21<sup>st</sup> century (Singhal *et al.*, 2008). It is due to its ability to store and yield more starch in its stem than that of any other starch crop (Onsa *et al.*, 2004). *Metroxylon sagu* is able to survive swamp area, acidic peat soils and in high saline area (Ehara *et al.*, 2000). It needs little management and able to live in marginal agricultural land (Purwanto *et al.*, 2002).

The plant is believed to have originated from Papua New Guinea and/or Moluccas where Papua New Guinea considered as the center of the diversity of sago palm (Toyoda, 2008). In Malaysia, Sarawak is the largest area for the *Metroxylon sagu* plantation which is approximately 55500 hectares in 2012 (Department of Agriculture Sarawak, 2013). Seventy-five percent of Sarawak sago planting area are found in Mukah (Bujang, 2010).



**Figure 1.1:** Sago plantation in Mukah, Sarawak (Bujang, 2008).

### **1.2.2 Sago processing**

Sago was processed commercially using the conventional and modern method in Sarawak (Shin and Collins, 2015). Conventional method and modern method had been described by Shin and Collins (2015) and Bujang (2008), respectively. Modern method usually being practiced in sago processing mills. In the modern method, mature sago trunk was selected and cut. After sago trunk is cut, it will be further cut into smaller one meter logs (Figure 1.2). The smaller logs of sago trunks will be transported to the sago processing mill by using lorries

or tied to form rafts so that they can be transported via river to the mill (Figure 1.3). Then, the bark of the sago trunk will be debarked manually or using automated debarking machine (Figure 1.4). The debarked logs are mashed using a rasper, followed by a hammer mill and then water is added to form starch slurry (Figure 1.5). Then, the starch slurry is allowed to settle down in the tank, dry and finally form sago flour that are ready for packaging (Figure 1.6). Figure 1.7 shows the schematic flow diagram of the sago processing.



**Figure 1.2:** Sago trunk was selected before cut into smaller 1 meter log.